



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

The Reply



We are aligned with Drs Olmos and Roque that there should be comprehensive medical crisis management as shown in the 4 pillars of pandemic response in the Figure.¹ Human severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection begins as a mild upper respiratory tract infection with a prehospital therapeutic window of opportunity, and there is an important “second pillar” of pandemic response that has the objective of reducing to the “hard outcomes” of COVID-19, hospitalization and death. For patients at high-risk of contracting coronavirus disease 2019 (COVID-19) and are acutely ill at home, contagion control is too late and the hospital is not an adequate safety net. Late-stage hospitalization as the only opportunity for initial treatment results in unacceptably high mortality.² Thus, in the context of a crisis, therapeutic decisions are made based on pathophysiological principles and the totality of available evidence. Because most serious viral infections require multidrug regimens, we can only expect

signals of efficacy or safety with single agents from randomized and observational studies of COVID-19. Clinical judgment is required to assemble therapeutic combinations that address viral replication, cytokine storm, and thrombosis.¹ Since the time of the original publication in *The American Journal of Medicine*, we are better supported from inpatient studies on the application of aspirin as well as anticoagulation that have established safety profiles.³ Meizlish et al⁴ found in a multicenter study ($N = 2785$) that aspirin administration was independently associated with a 69% reduction in mortality ($P = .001$). Billet et al⁵ ($N = 3625$) demonstrated a significant decrease in adjusted mortality with prophylactic use of apixaban (odds ratio = 0.46, $P = .001$) and enoxaparin (odds ratio = 0.49, $P = .001$). We encourage Drs Olmos and Roque to overcome the fear of relying on clinical judgment before confirmatory large-scale multidrug, placebo-controlled, randomized trials. To our knowledge no such trials are forthcoming. Empiric regimens based on clinical judgment are not as “dangerous” as leaving patients untreated for many days only to succumb to calamitous hospitalization



Figure The 4 pillars of pandemic response to the severe acute respiratory coronavirus 2 (SARS-CoV-2) pandemic crisis, reproduced with permission from McCullough et al.¹

Funding: None.

Conflicts of Interest: None.

Authorship: The author is solely responsible for the content of this manuscript.

Requests for reprints should be addressed to Peter A. McCullough, MD, MPH, Baylor Heart and Vascular Institute, 621 N. Hall St, H030, Dallas, TX, 75226.

E-mail address: peteramccullough@gmail.com

or death. Courageous doctors and researchers have innovated and found that sequenced, multidrug regimens are associated with ~85% reductions in COVID-19 morbidity and mortality with no signals of harm.^{6,7,8}

Peter A. McCullough, MD, MPH^{a,b,c}

^a*Baylor University Medical Center,
Dallas, Tex*

^b*Baylor Heart and Vascular
Institute, Dallas, Tex*

^c*Baylor Jack and Jane Hamilton
Heart and Vascular Hospital,
Dallas, Tex*

<https://doi.org/10.1016/j.amjmed.2021.01.013>

References

1. McCullough PA, Alexander PE, Armstrong R, et al. Multifaceted highly targeted sequential multidrug treatment of early ambulatory high-risk SARS-CoV-2 infection (COVID-19). *Rev Cardiovasc Med* 2020;21(4):517–30. <https://doi.org/10.31083/j.rcm.2020.04.264>.
2. Palazzuoli A, Ruberto F, De Ferrari GM, et al. Inpatient mortality according to level of respiratory support received for severe acute respiratory syndrome coronavirus 2 (Coronavirus Disease 2019) infection: a prospective multicenter study. *Crit Care Explor* 2020;2(9): e0220. <https://doi.org/10.1097/CCE.0000000000000220>.
3. McCullough PA, Kelly RJ, Ruocco G, et al. Pathophysiological basis and rationale for early outpatient treatment of SARS-CoV-2 (COVID-19) infection. *Am J Med* 2021;134(1):16–22. <https://doi.org/10.1016/j.amjmed.2020.07.003>.
4. Meizlish ML, Goshua G, Liu Y, Fine R, et al. Intermediate-dose anticoagulation, aspirin, and in-hospital mortality in COVID-19: a propensity score-matched analysis [e-pub ahead of print]. medRxiv. Accessed December 30, 2020. doi: <https://doi.org/10.1101/2021.01.12.21249577>.
5. Billett HH, Reyes-Gil M, Szymanski J, et al. Anticoagulation in COVID-19: effect of enoxaparin, heparin, and apixaban on mortality. *Thromb Haemost* 2020;120(12):1691–9. <https://doi.org/10.1055/s-0040-1720978>.
6. Procter BC, Ross C, Pickard V, Smith E, Hanson C, McCullough PA. Early ambulatory multidrug therapy reduces hospitalization and death in high-risk patients with SARS-CoV-2 (COVID-19) [e-pub ahead of print]. *Authorea*. Accessed January 25, 2021. doi: <https://doi.org/10.22541/au.161000355.54720791/v1>.
7. Procter BC, Ross C, Pickard V, Smith E, Hanson C, McCullough PA. Clinical outcomes after early ambulatory multidrug therapy for high-risk SARS-CoV-2 (COVID-19) infection. *Rev Cardiovasc Med* 2020;21(4):611–4. <https://doi.org/10.31083/j.rcm.2020.04.260>.
8. Derwand R, Scholz M, Zelenko V. COVID-19 outpatients: early risk-stratified treatment with zinc plus low-dose hydroxychloroquine and azithromycin: a retrospective case series study. *Int J Antimicrob Agents* 2020;56(6):106214. <https://doi.org/10.1016/j.ijantimicag.2020.106214>.